

GENETIC VARIABILITY OF microRNA GENES IN FARM ANIMALS

Daša JEVŠINEK SKOK ¹, Minja ZORC ^{1,2}, Simon HORVAT ^{1,3}, Peter DOVČ ¹, Milena KOVAČ ¹,
Tanja KUNEJ ¹

ABSTRACT

MicroRNAs (miRNAs) are a class of non-coding RNAs that plays an important role in posttranscriptional regulation of target genes. Regulation requires complementarity between the target mRNA and the miRNA seed region, which is responsible for their recognition and binding. Previous studies in human and mouse have shown that variability of miRNA gene (miR-SNPs) might interfere with its function resulting in phenotypic variation. Polymorphisms within miRNA genes could represent biomarkers for phenotypic traits important in farm animals. The aim of this study was to: 1) update previously developed web-based tool for identification of polymorphisms within miRNA genes (miRNA SNIPer), 2) systematically collect polymorphisms of miRNA genes in pig, cattle, chicken, and horse, and 3) experimentally validate SNPs within miRNA seed regions (miR-seed-SNPs) in cattle. Using miRNA SNIPer tool, polymorphisms within 32 mature miRNA regions, including 12 miR-seed-SNPs, were identified in pig, cattle, and chicken. Bovine miR-seed SNPs were chosen for experimental validation. The bta-mir-2313 locus was shown to be very polymorphic, therefore we validated one SNP with previously unknown validation status within the mature seed region in population of Slovenian Simmental cattle. Additionally, two SNPs in corresponding pri-miRNA were identified. Results of this study can serve researchers for follow up hypothesis-driven experimental studies to evaluate the phenotypic effect of identified miRNA genetic variability in vertebrates.

Keywords: farm animals / biomarkers / microRNA / genetic variability

1 INTRODUCTION

MicroRNAs (miRNAs) are non-coding RNA molecules with approximately 21 nucleotides in length that play an important role in posttranscriptional regulation of mRNA. By binding to the different target gene regions, *i.e.*, 3' untranslated region (3'UTR), 5'UTR, promoter, or coding sequences, they repress or activate translation (reviewed in Kunej *et al.*, 2012). MicroRNA biogenesis begins in the nucleus with the primary transcript (pri-miRNA) of several hundreds or thousands base pairs in length that is cleaved by the action of endonuclease DROSHA to 60 to 70 nucleotides long precursor miRNA (pre-miRNA). Pre-miRNA, with its characteristic stem-loop structure (Fig. 1) is then transported to the cytoplasm

where endonuclease DICER cleaves both duplex chains to form mature miRNA (Lee *et al.*, 2002; Bartel, 2004). Products of DICER action also include complementary sequences of mature miRNAs, which are referred to as miRNA* (Lau *et al.*, 2001), and are usually transcribed in lower percentage as mature miRNAs (Lim *et al.*, 2003). The key binding location for translational suppression, also called the seed region, resides in the mature miRNA sequence, more accurately situated at position 2–7 or 2–8 nucleotides from the 5' end of the miRNA (Sun *et al.*, 2009).

Changes in the miRNA expression profile were linked with several diseases (reviewed in Ferdin *et al.*, 2011). Moreover, single nucleotide polymorphisms (SNPs) within 1) miRNA genes, 2) miRNA targets or in

¹ Univ. of Ljubljana, Biotechnical Fac., Dept. of Animal Science, Groblje 3, SI-1230 Domžale, Slovenia

² Centre for Mathematical and Computational Biology, Rothamsted Research, Harpenden, UK

³ Dept. of Biotechnology, National Institute of Chemistry, Ljubljana 1000, Slovenia




Figure 1: Secondary structure of miRNA and an example of miRNA stem-loop (adapted from Saunders *et al.*, 2007)

3) protein-coding genes involved in miRNA biogenesis result in phenotypic differences and therefore affects (associated with) production traits and susceptibility to diseases (Georges *et al.*, 2007). Previously, we developed a web based tool miRNA SNIper for the identification of polymorphisms residing within miRNA genes (Zorc *et al.*, 2012). Since some farm animals (pig, cattle, chicken,

sheep, etc.) represent both – a source of food and disease models, the biomarkers based on miRNA polymorphisms could contribute to the study of phenotypic properties in farm and in medicine. In this study we systematically collected polymorphic miRNA genes from four farm animal species (pig, cattle, chicken, and horse) and further experimentally validated miRNA SNPs in cattle.

miRNA SNIper

Tool for identification of genetic variations residing within seed microRNA regions, responsible for mRNA binding in vertebrates: Human (eg. *hsa-mir-941-3*), Mouse (eg. *mmu-mir-654*), Cattle (eg. *bta-mir-29e*)

Enter microRNA names:

bta-mir-2901
bta-mir-2902-1
bta-mir-2902-2
bta-mir-2903
bta-mir-2904-1
bta-mir-2904-2
bta-mir-2904-3
bta-mir-2917
bta-mir-2957
bta-mir-3431
bta-mir-3432-1
bta-mir-3432-2
bta-mir-3578
bta-mir-3596
bta-mir-3600
bta-mir-3601
bta-mir-3602
bta-mir-3604-1
bta-mir-3604-2

Send

Results:

bta-mir-2313

Species: *Bos taurus*

Mirna location: 15:32789445-32789518 Strand:+

Seed location: 15:32789495-32789501 Strand:+

GGGCUGGAGUGCAGCUGAGGACCAAGGCAGGGCUGCAUGCAUUCA
CAUGCCAGUCCACGCUGCAUGCGGCCU

Premature miRNA variation rs41761413

SNV

SNP location: 32789501

Reference sequence: C

Variant sequence: T

seed SNP rs41761413

SNV on 7. nucleotid of seed.

SNP location: 32789501

Reference sequence: C

Variant sequence: T

Figure 2: An example of predicted genetic variability within miRNA in cattle (*bta-mir-2313*), using a web-based tool miRNA SNIper. Gray: mature miRNA, underlined region: seed region

2 MATERIALS AND METHODS

2.1 UPDATE OF THE ONLINE TOOL FOR THE DETECTION OF GENETIC VARIATIONS WITHIN MIRNA GENES.

Previously developed tool miRNA SNiPer was updated with the latest versions of miRBase (release 18; <http://www.mirbase.org/>) (Kozomara and Griffiths-Jones, 2011), TargetScan (release 5.2; <http://www.targetscan.org/>) (Lewis *et al.*, 2005), and Ensembl Variation database for pig (Sscrof10.2), horse (EquCab2), cattle (Btau4.0), and chicken (WASHUC2).

2.2 SAMPLES, SEQUENCING OF A DNA PANEL OF SIRES AND VALIDATION OF MIR-SNPS IN CATTLE

Animals were selected from the National progeny test for Slovenian Simmental cattle. DNA samples were extracted from frozen semen of sires using DNeasy Blood & Tissue DNA extraction kit (QiaGen, Düseldorf, Germany). Three bovine miR-seed-SNPs were chosen for experimental validation. Primers were selected using online tool Primer3 (Rozen and Skaletsky, 2000). Due to presence of the *repetitive sequence*, primers in region of SNP rs42658514 within bta-mir-2450c couldn't be designed. For experimental validation, the following primers were used: for bta-mir-2313 forward primer (F) 5'-GCACA-

GACTCTCAGCCACTG-3' and reverse primer (R) 5'-CTGACTGAGGCTCTCGCTCT-3', and for bta-mir-29e (F) 5'-TGTAGGGACTGGTTGTGGAA-3' and (R) 5'-TCTACTGAACACAGCCCCATC-3'. PCR products were purified using *ExoI* (Exonuclease I) and SAP (shrimp alkaline phosphatase (both Fermentas, Vilnius, Lithuania) and following sequencing reaction for capillary electrophoresis on ABI3130xl.

3 RESULTS AND DISCUSSION

Previously developed web-based tool miRNA SNiPer, designed for the detection of genetic variations within miRNA genes in vertebrates, was updated. The tool accepts a list of miRNA genes and returns a table of variations within different regions of miRNA genes: pre-miRNA, mature, and seed region (Fig. 2).

The number of total known miRNAs and putative SNPs, as well as the list of miR-SNPs for four farm animal species (pig, horse, cattle, and chicken) is presented in table 1. The highest number of SNPs in farm animals is currently known for chicken and cattle, while the numbers for other farm animals are significantly lower. Thirty-two polymorphisms overlapping mature miRNAs, including 12 within seed region (miR-seed-SNPs), were identified. One miR-seed-SNP was identified in pig, three in cattle, and seven in chicken. Bovine miR-seed-SNPs were selected for experimental validation in eight Slovenian Simmental cattle sires.

All collected miR-seed-SNPs have an unknown

Table 1: Number of known miRNAs and SNPs within four farm species (pig, cattle, chicken, and horse) and list of SNPs within miRNA mature and seed region

| Species | Total number of known miRNAs | Total number of SNPs | No. of SNPs within seed/mature miRNA region | miRNA comprising seed SNP | miR-seed-SNP ID |
|---------|------------------------------|----------------------|---|--|--|
| Pig | 228 | 545.950 | 1/8 | ssc-mir-4335 | rs80984906 |
| Cattle | 662 | 2.201.071 | 3/9 | bta-mir-29e bta-mir-2313 bta-mir-2450c | rs41825418 rs41761413 rs42658514 |
| Chicken | 499 | 3.292.991 | 8/14 | gga-miR-1568 gga-miR-1614* gga-miR-1644 gga-miR-1648* gga-miR-1657 gga-miR-1658 | rs14511527 rs15172520 rs14076349 rs14281065 rs14934924 rs16681031 rs16681032 rs16681033 |
| Horse | 341 | 1.163.258 | 0/1 | / | / |




Figure 3: Validation of three pri-miRNA SNPs located within bta-mir-2313 in cattle. Gray: mature miRNA, underlined region: seed region

validation status according to the NCBI database and have not been genotyped yet. Experimental validation revealed that polymorphism rs41825418 within bta-mir-29e was monomorphic, while miR-seed-SNP rs41761413 within bta-mir-2313 was polymorphic in analyzed population of Slovenian Simmental cattle (Fig. 3). Additional two SNPs in corresponding pri-miRNA were identified: rs41761412 with poly-allelic substitution (G> T> A) and rs41761414 with substitution (A> G).

4 CONCLUSION

Since miRNA polymorphisms may have a profound effect on a wide range of phenotypes, genetic variability of miRNA genes in farm animals was examined. Bioinformatics tool miRNA SNiPer was updated and used for assembling a list of all known miRNA SNPs in four farm species (pig, horse, cattle, and chicken). Our results show, that most of the miR-SNPs still need to be validated. The project is ongoing, as novel miRNAs and SNPs are yet to be discovered in farm animals, as well as in other animal species. Collated data can be used by interested scientific community to help retrieve valuable information and design efficient experimental plans in the field of miR-

SNP research. Association study between validated SNPs located within bta-mir-2313 and carcass traits in paternal half-sib families of the Slovenian Simmental cattle is still under way and current results will be presented. This project may yield new findings useful for development of molecular markers in selection programs allowing more effective, marker assisted selection in farm animals.

5 REFERENCES

- Bartel D.P. 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell*, 116: 281–297
- Ferdin J., Kunej T., Calin G.A. 2011. MicroRNA: Genomic Association with Cancer Predisposition. *Journal of the Association of Genetic Technologists*, 37: 11–19
- Georges M., Coppieters W., Charlier C. 2007. Polymorphic miRNA-mediated gene regulation: contribution to phenotypic variation and disease. *Current Opinion in Genetics and Development*, 17: 166–176
- Kozomara A., Griffiths-Jones S. 2011. miRBase: integrating microRNA annotation and deep-sequencing data. *Nucleic Acids Research*, 39: D152–D157
- Kunej T., Godnic I., Horvat S., Zorc M., Calin G.A. 2012. Cross Talk Between MicroRNA and Coding Cancer Genes. *The Cancer Journal*, 18: 223–231
- Lau N.C., Lim L.P., Weinstein E.G., Bartel D.P. 2001. An abun-

- dant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science*, 294: 858–862
- Lee Y., Jeon K., Lee J.-T., Kim S., Kim V.N. 2002. MicroRNA maturation: stepwise processing and subcellular localization. *The EMBO Journal*, 21: 4663–4670
- Lewis B.P., Burge C.B., Bartel D.P. 2005. Conserved Seed Pairing, Often Flanked by Adenosines, Indicates that Thousands of Human Genes are MicroRNA Targets. *Cell*, 120: 15–20
- Lim L.P., Lau N.C., Weinstein E.G., Abdelhakim A., Yekta S., Rhoades M.W., Burge C.B., Bartel D.P. 2003. The microRNAs of *Caenorhabditis elegans*. *Genes and Development*, 17: 991–1008
- Rozen S., Skaletsky H. 2000. Primer3 on the WWW for general users and for biologist programmers. *Methods in Molecular Biology*, 132: 365–386
- Saunders M.A., Liang H., Li W.H. 2007. Human polymorphism at microRNAs and microRNA target sites. *Proceedings of the National Academy of Sciences of the United States of America*, 104: 3300–3305
- Sun G., Yan J., Noltner K., Feng J., Li H., Sarkis D.A., Sommer S.S., Rossi J.J. 2009. SNPs in human miRNA genes affect biogenesis and function. *RNA*, 15: 1640–1651
- Zorc M., Jevsinek Skok D., Godnic I., Calin G.A., Horvat S., Jiang Z., Dovc P., Kunej T. 2012. Catalog of MicroRNA Seed Polymorphisms in Vertebrates. *PloS one*, 7: e30737